

EDITORIAL

ANCA-positive Vasculitides and Coronary Artery Pathology

Haralambos M. Moutsopoulos, MD

Department of Pathophysiology,
School of Medicine, National
University of Athens, Athens, Greece

KEY WORDS: *vasculitis; coronary ectasia; magnetic resonance imaging*

LIST OF ABBREVIATIONS:

ANCA = antibodies to neutrophil cytoplasmic antigens
ceMRI = contrast-enhanced MRI
CS = Churg-Strauss syndrome
Gd-DTPA = gadolinium-diethylene triamine pentaacetic acid
LAD = left anterior descending (coronary artery)
LCx = left circumflex
MRA = magnetic resonance angiography
MPA = microscopic polyangiitis
PAN = polyarteritis nodosa
RA = rheumatoid arthritis
RCA = right coronary artery
WG = Wegener's granulomatosis

Correspondence to:

Haralambos M. Moutsopoulos, MD,
FACP, FRCP (Edin)
Professor and Director
Department of Pathophysiology
School of Medicine
National University of Athens
75 Mikras Asias Street
115 27 Athens, Greece
Tel.: + 30 210 746 2512 - 14
Fax.: + 30 210 746 2664
E-mail.: hmoutsop@med.uoa.gr

ABSTRACT

Systemic necrotizing vasculitides preferentially involve small- or medium-sized vessels and occasionally the coronary vessels and the myocardium. Cardiac involvement was noninvasively evaluated in a group of 32 patients with vasculitides but without cardiac symptoms with use of magnetic resonance imaging techniques and compared with a control group of 13 individuals. The study showed increased coronary vessel diameters in the majority of patients and in some patients to the degree of ectasia; myocardial scar was rare.

Vasculitides constitute a heterogeneous group of disorders affecting from large arteries to post-arteriolar capillaries. These disorders can affect all organs and are categorized on the basis of the affected artery diameter and the presence or absence of antibodies to neutrophil cytoplasmic antigens (ANCA). ANCA-positive vasculitides involve small to medium size arteries of many organs, primarily of the upper and lower respiratory system, the kidneys and the peripheral nervous system. Microscopic polyangiitis (MPA), Wegener granulomatosis (WG) and Churg-Strauss syndrome (CS) are forms of these vasculitides.¹ Coronary arteries have been previously shown to be occasionally affected during the course of these diseases; however, this has never been formally assessed.

To evaluate non-invasively the coronary arteries of patients with ANCA-positive vasculitides, magnetic resonance angiography (MRA) was performed and the viability was studied using contrast-enhanced MRI (ceMRI).² Twelve patients with MPA, 13 with WG, 5 with CS and 2 with polyarteritis nodosa (PAN), all without any cardiac symptoms were included in the study and compared with 13 age- and sex-matched healthy controls and with a disease control group of 13 patients with rheumatoid arthritis (RA). The maximal diameter of the proximal one third of each coronary vessel was recorded. MRA was performed using a 1.5 Tesla system. Data acquisition was performed with ECG gating in mid-diastole and with the patient breathing freely. ceMRI images were acquired 15 minutes after the IV injection of 0.15 mmol/kg gadolinium-diethylene triamine pentaacetic acid (Gd-DTPA) using an inversion recovery gradient echo pulse sequence.

As shown in the table (Table 1), the diameter of all coronary arteries tested were found significantly increased in patients with MPA, PAN and WG ($p < 0.001$ for LAD

TABLE 1. Coronary vessel diameters (mean \pm SD) in patients and controls

Coronary vessel	MPA+PAN *	WG *	CS	RA	Healthy controls
LAD (mm)	4.58 \pm 1.27	3.57 \pm 0.69	2.66 \pm 0.51	3.06 \pm 0.60	3.05 \pm 0.15
RCA (mm)	4.68 \pm 1.00	3.55 \pm 0.92	2.68 \pm 0.77	3.27 \pm 0.81	3.17 \pm 0.20
LCx (mm)	4.01 \pm 1.16	3.16 \pm 0.45	2.48 \pm 0.73	3.20 \pm 0.56	3.08 \pm 0.19

* P <0.01 vs controls

CS= Churg-Strauss syndrome; LAD= left anterior descending (coronary artery); LCx= left circumflex; MPA= microscopic polyangiitis; PAN= polyarteritis nodosa; RA= rheumatoid arthritis; RCA= right coronary artery; WG= Wegener's granulomatosis

and RCA, $p < 0.01$ for LCx) compared to controls, but not in those with CS. Criteria for ectasia were fulfilled by MPA and PAN patients, but not WG. The comparison of coronary vessel diameters revealed increased coronary artery diameter in the MPA+PAN patient group compared to that of WG ($p < 0.05$ for LAD and LCx, $p < 0.01$ for RCA). Coronary vessel diameters were increased in the MPA+PAN group compared to RA ($p < 0.01$ for LAD and RCA and $p < 0.05$ for LCx), but there was no difference between WG or CS and RA. Evidence of myocardial scar in Gd-DTPA images was documented in 1 patient with MPA and in 1 patient with CS.

In **conclusion**, coronary artery ectasia appears a common finding in asymptomatic vasculitis patients, while scar is rare. Magnetic resonance imaging³ evaluation is feasible in these

forms of vasculitis and may prove of value for treatment guidance.

REFERENCES

1. Jennette JC, Wilkman AS, Falk RJ. Diagnostic predictive value of ANCA serology. *Kidney Int* 1998;53:796-798.
2. Mavrogeni S, Manoussakis MN, Karagiorga TC, et al. Detection of coronary artery lesions and myocardial necrosis by magnetic resonance in systemic necrotizing vasculitides. *Arthritis Rheum* (in press).
3. Botnar RM, Stuber M, Danias PG, Kissinger KV, Manning WJ. Improved coronary artery definition with T2-weighted, free breathing three-dimensional coronary MRA. *Circulation* 1999;99:3139-3148.